

Global Embedding of Analytic Branes into $\mathcal{E}^\Sigma$ Einstein MD Bulk Cosmology: Wormholes

Nikolaos I. Katzourakis*

University of Athens, Department of Mathematics

e-mail: nkatzourakis@math.uoa.gr

Abstract

In a series of recent papers [26], [27], [28] we have shown that:

(I) *any n -dimensional analytic semi-Riemannian manifold $\mathcal{M}$ has a global isometric analytic embedding into an Einstein manifold $\mathcal{E}$ of dimension $n + d$, $d \geq 1$ ($\Lambda \in \mathbb{R}$).*

(II) *There exists an Einstein global embedding space $\mathcal{E}^\Sigma$, admitting unsmoothable differential-topological singularities ($0 - D$ points or extended $(d - k) - D$ spatial "anomalies") living into the complement $\mathcal{E}^\Sigma \setminus \mathcal{M}$. They were expounded as wormholes.*

(III) *Any finite-many analytic semi-Riemannian manifolds $\mathcal{M}_1^{(n_1)}, \dots, \mathcal{M}_m^{(n_m)}$ have all a global isometric analytic embedding into an $(\sum_{1 \leq k \leq m} n_k + d)$ -Einstein Bulk $\mathcal{E}$ as disjoint submanifolds, while $\pi_m(\mathcal{E})$ splits. The optimum dimensionality corresponds to $d = 1$.*

In this paper we continue extending further these results on Cosmological Brane embeddings:

There exists an Einstein global embedding space $\mathcal{E}^\Sigma$, admitting unsmoothable differential-topological singularities ($0 - D$ points or extended $(d - k) - D$ spatial "anomalies") living into the complement $\mathcal{E}^\Sigma \setminus \prod_{k=1}^m \mathcal{M}_k$.

*Address: Troias 6, 18541, Kaminia, Pireaus, Greece

Keywords: Spacetime Embedding, Einstein space, Campbell-Magaard Theorem; Brane-World, MD Gravity, Brane Cosmology

MSC Subject Classification 2000: Primary 53C20, 53C25, 32C09, 53Cxx; Secondary 83C75, 83C55, 83E15, 83F05.

Introduction

The concept of the *spacetime manifold* as a model of the large scale structure of the Universe is a result of evolution in both physical & mathematical attitudes. Once we used to think of Euclidean flat 3-space as the case of interest in mathematics, constituting in the same time the model of the Newtonian space-time.

In the first half of the 20th century, Einstein's General Relativity (*G.T.R.*) employed the differential-geometric model of 4-*manifold* equipped with a *Lorentzian metric* structure, the solution of the *field equations* which (the latter) correlate the *Curvature* of spacetime with the distribution of *matter*, the *energy* and the *momentum* ([5], [7], [12], [14]). That was the beginning of interconnections between geometry & physics.

The possibility that there may exist more than $3 + 1$ space-time dimensions was taken into account not long after the final formulation of G.T.R.. *Kaluza* and *Klein* (K.-K.) *unified 4D vacuum Relativity* and classical *electromagnetism* ([15]) on a 5-*manifold* that satisfies the vacuum field equations (curved by the contribution of the photons). The extra dimension was considered topologically *compact* (*homeomorphic to $\mathbb{S}^1$*), *infinitesimal* (thus invisible) and the 4-spacetime was assumed not to depend on the 5th dimension. In fact, the Universe was considered as a *hyper-cylinder*, locally *homeomorphic to $\mathbb{R}^4 \times \mathbb{S}^1$* .

In a modern approach called *Space-Time-Matter* or *Induced Matter Theory* (S.T.M.) ([17], [18], [15] and references therein) these assumptions are removed and the result is the *unification of G.T.R. & electromagnetism together with their sources* (*mass & charge*), expounded (the latter) as an artifact induced on the 4-manifold, generated by the (non-linear) geometry of invisible extra spatial dimensions.

One of the most recent Cosmological scenarios concerning the geometric struc-

ture of the (classical) Universe introduced by L. Randall and R. Sundrum ([38], [39]) claims that the Universe of our perception is a (mem-)Brane of $3 + 1$ dimensions (submanifold) embedded in a multi-dimensional (MD) Bulk of at least 5 dimensions. The main ansatz for this (possibly moving in the Bulk) Brane is that *matter is confined on the Brane and only gravity and electromagnetic fields can propagate on the bulk*. Soon, 2-Brane ($\mathcal{E}$ multi-Brane, colliding or not) systems embedded into the same Bulk came into light, with possible MD singularities. Our visible Universe was thought of as 1 of those interacting Branes.

Mathematically, these theories are modeled on a local embedding theorem of Riemannian Geometry, due to Campbell who was the first who stated it [2], and Magaard ([8], [31]) who presented a strict proof (see [35], [40], [44]): *any analytic n -manifold can be locally and isometrically embedded into some other Ricci-flat of $n+1$ dimensions ($R_{AB} = 0$)*. In a sense, this is equivalent to the *local embedding of Einstein 4D Relativity, into some 5D Relativity in the vacuum where no matter exists* ("wood", according to Einstein), but only *geometrical properties* ("marble").

It becomes clear that in the light of these theories that are entirely based on differential-geometric (though *local* only) embeddings, new *global results are needed*, in order to provide a true *supporting mathematical framework for Brane-Worlds*, not depending on local coordinates. The necessity for embedding owes to the fact that the starting point of an MD gravitational-cosmological theory is the 4-dimensional G.T.R. that has to be incorporated in the new MD context.

In recent work ([26], [27], [28]) the author has proved that **the Campbell-Magaard (C.-M.) Theorem** (extended by *Dahia & Romero* in [17] on *Einstein from Ricci-flat spaces*) **holds true globally**:

Any n -dimensional analytic semi-Riemannian manifold $\mathcal{M}$ has a global isometric analytic embedding into an Einstein manifold $\mathcal{E}$ of dimension $n + d$, $d \geq 1$ ($\Lambda \in \mathbb{R}$).

This $\mathcal{E}$ may admit (if desired) countable unsmoothable differential-topological singularities, which may be distinct points into spacetime or extended spatial anomalies ([27]). These pathologies living into the complement $\mathcal{E} \setminus \mathcal{M}$ were expounded as wormholes connecting distant points of spacetime.

In addition, Any analytic semi-Riemannian manifolds $\mathcal{M}_1^{(n_1)}, \dots, \mathcal{M}_m^{(n_m)}$ have all a global isometric analytic embedding into an $(\sum_{1 \leq k \leq m} n_k + d)$ -Einstein Bulk $\mathcal{E}$ as disjoint submanifolds, while both $\pi_m(\mathcal{E})$ & $H_m(\mathcal{E}; \mathbb{G})$ split. The optimum dimensionality of $\mathcal{E}$ corresponds to embedding codimension $d = 1$.

In this paper we combine certain ideas from the prequels, showing that the Einstein embedding manifold $\mathcal{E}$ of the Branes may admit 1 in codimension $d = 1$ & countable-many in $d > 1$ unsmoothable differential-topological singularities, which may be distinct points into spacetime or extended spatial anomalies.

The procedure in the proofs is a slight modification of the one we followed in [26] & [27]. We form a bulk that contains all the $\mathcal{M}_k$'s as disjoint submanifolds and using the known local embedding results and certain differential-geometric & topological arguments we obtain the requested global extension.

In *Section 1* we state the results proved in the prequels. *Section 2* contains the proof of the basic result stated above. In *Section 3* we study the topology of such embedding Einstein spaces, giving a Homological description of the Einstein manifolds $\mathcal{E}$ we construct, illustrating the aforementioned splitting.

The analogous physical interpretation of the constructed *pathological anomalies as wormholes connecting distant points of Bulk spacetime* living in the extra dimensions holds in this case of multi-Brane systems too. We refer to [27].

Notations: In the sequel, the terms **Spacetime**, **Brane**, **Bulk** will be used in the place of *paracompact*, *Hausdorff*, *connected manifolds (finite dimensional)*, equipped with a *semi-Riemannian metric structure* and the *corresponding Levi-Civita linear connection*.

1. The Previous Results.

In [26] we proved that:

Theorem 1.1. (*Global Isometric Embedding into Einstein manifolds in codimension 1*)

Any n -dimensional real analytic semi-Riemannian manifold $(\mathcal{M}; \nabla^{\mathcal{M}}, g_{\mathcal{M}})$ has a global isometric embedding into an $(n + 1)$ -dimensional Einstein manifold

$(\mathcal{E}; \nabla^\mathcal{E}, g_\mathcal{E})$ satisfying:

$$Ric^{\nabla^\mathcal{E}}(g_\mathcal{E}) = \frac{2\Lambda}{n-1} g_\mathcal{E}$$

for any $\Lambda \in \mathbb{R}$.

It is worth noticing that the assumption of analyticity for the manifold is not restrictive, provided that *any paracompact C^k -manifold, $0 \leq k \leq \infty$, acquires a unique compatible not just smooth (C^∞) but real analytic structure* (there exists a C^k -diffeomorphism with an analytic manifold). See [45], [24], [10].

Remark 1.2. It is obvious that if a manifold can be embedded into an Einstein space as a hypersurface, repeated applications of *Theorem 1.1* imply the weaker condition to be *embedable in any codimension greater than 1*.

In [27] we extended further this result, so that to incorporate pathological singularities in the complement of the Bulk $\mathcal{E} \setminus \mathcal{M}$ that are non-removable:

Theorem 1.3. (*Embedding into Einstein manifold with unsmoothable differential-topological singularities of zero measure*)

Given a semi-Riemannian analytic manifold $\mathcal{M}$, there exist:

- (I) a topological space $(\mathcal{E}^\Sigma, \mathcal{T}_{\mathcal{E}^\Sigma})$
- (II) an analytic manifold $(\mathcal{E}, \mathcal{A}_\mathcal{E})$, with $\dim(\mathcal{E}) = \dim(\mathcal{M}) + d$, $d \geq 1$
- (III) a countable set of points $\mathbb{F}^\Sigma \subseteq \mathcal{E}^\Sigma$, satisfying:

$$\mathbb{F}^\Sigma = \begin{cases} \{f^\sigma\}, & \text{if } d = 1 \\ \{f_{(1)}^\sigma, f_{(2)}^\sigma, \dots\} \cong \mathbb{N}, & \text{if } d \geq 1. \end{cases}$$

such that the following topological homeomorphism holds (under inclusion to a subspace)

$$(\mathcal{E}^\Sigma \setminus \mathbb{F}^\Sigma, \mathcal{T}_{\mathcal{E}^\Sigma}|_{\mathcal{E}^\Sigma \setminus \mathbb{F}^\Sigma}) \xrightarrow[\supset]{\cong} (\mathcal{E}, \mathcal{T}_{\mathcal{A}_\mathcal{E}})$$

and $\mathcal{E}$ is an Einstein space of global isometric analytic embedding for $\mathcal{M}$.

In addition, $\mathcal{E}^\Sigma$ is **not** even a topological manifold and there does not exists any smoothing resolution of the singularities $\mathbb{F}^\Sigma \subseteq \mathcal{E}^\Sigma$.

$\mathcal{T}_{\mathcal{A}_\mathcal{E}}$ denotes the (induced from the differential structure) topology of $\mathcal{E}$.

The mathematical mechanism of topological identification we used in the proof of 1.3 in [27] to produce the pathological singularities on the Bulk raised the idea that it is not really necessary these "anomalies" to be points: there may exist extended problematic domains of the complement $\mathcal{E}^\Sigma \setminus \mathcal{M}$. We simply identify distinct homeomorphic domains on the "fibres" $\mathcal{F}$.

This more realistic case may correspond to extended object in the Universe with positive (multi-dimensional) volume.

Theorem 1.4. (*Embedding into Einstein manifold with unsmoothable differential-topological singularities of positive $(d - k)$ -measure*)

Given a semi-Riemannian analytic manifold $\mathcal{M}$, there exist:

- (I) *a topological space $(\mathcal{E}^\Sigma, \mathcal{T}_{\mathcal{E}^\Sigma})$*
- (II) *an analytic manifold $(\mathcal{E}, \mathcal{A}_\mathcal{E})$, with $\dim(\mathcal{E}) = \dim(\mathcal{M}) + d$, $d \geq 1$*
- (III) *countable many closed path connected $\mathcal{C}$ simply connected sets $\mathbb{F}^\Sigma \subseteq \mathcal{E}^\Sigma$,*

satisfying:

$$\mathbb{F}^\Sigma = \begin{cases} \{f^\Sigma\} \cong [0, 1], & \text{if } d = 1 \\ \{f_{(1)}^\Sigma, f_{(2)}^\Sigma, \dots\} \cong \mathbb{N} \times \overline{\mathbb{B}^{d-k}(0, 1)}, & \text{if } d \geq 2. \end{cases}$$

($0 \leq k \leq d$) such that the following topological homeomorphism holds (under inclusion to a subspace)

$$(\mathcal{E}^\Sigma \setminus \mathbb{F}^\Sigma, \mathcal{T}_{\mathcal{E}^\Sigma}|_{\mathcal{E}^\Sigma \setminus \mathbb{F}^\Sigma}) \xrightarrow[\supset]{\cong} (\mathcal{E}, \mathcal{T}_{\mathcal{A}_\mathcal{E}})$$

and $\mathcal{E}$ is an Einstein space of global analytic isometric embedding for $\mathcal{M}$.

*In addition, $\mathcal{E}^\Sigma$ is **not** a topological manifold.*

In [28] we showed that there *exists a supporting embedding result* for the *multi-Brane models* of Cosmology (see e.g. [42]), due to the following result.

The method in the proof is a slight modification of the 1-manifold embedding of [26] to produce an $\mathcal{E}$ admitting finite many submanifolds.

Theorem 1.5. (*Global Isometric Embedding of Branes into Einstein manifolds in codimension 1*)

Any n_k -dimensional analytic semi-Riemannian manifolds $(\mathcal{M}_k; \nabla^{\mathcal{M}_k}, g_{\mathcal{M}_k})$, $1 \leq k \leq m$ (Branes) have all a global analytic isometric embedding into an

$(\sum_{k=1}^m n_k + 1)$ -dimensional Einstein manifold $(\mathcal{E}; \nabla^\mathcal{E}, g_\mathcal{E})$ satisfying:

$$Ric^{\nabla^\mathcal{E}}(g_\mathcal{E}) = \frac{2\Lambda}{\left(\sum_{k=1}^m n_k\right) - 1} g_\mathcal{E}$$

as disjoint submanifolds for any $\Lambda \in \mathbb{R}$.

In this paper we continue with the reasonable extension of the last theorem concerning Brane embeddings incorporating singular spacetimes, in the sense in was done in [27] in the case of a single manifold embedding.

2. Embedding of Branes into $\mathcal{E}^\Sigma$ C_α^∞ -Manifolds admitting Singularities.

The singularities we produce on $\mathcal{E}^\Sigma$ are *by construction pathological* and there does **not** exist *any resolution* of them: they are obtained by certain topological identifications of distinct point and the obtain Bulk is not even a manifold, unless the singularities are extracted.

In other words, there exists an Einstein Bulk $\mathcal{E}$ imbedded (in the sense of analysis) into a topological space $\mathcal{E}^\Sigma$, where the complement $\mathcal{E}^\Sigma \setminus \mathcal{E}$ has zero measure.

Theorem 2.1. (*Embedding into Einstein manifold with unsmoothable differential-topological singularities of zero measure*)

Given any analytic semi-Riemannian manifolds (Branes) $\mathcal{M}_1, \dots, \mathcal{M}_m$ there exist:

- (I) a topological space $(\mathcal{E}^\Sigma, \mathcal{T}_{\mathcal{E}^\Sigma})$
- (II) an analytic manifold $(\mathcal{E}, \mathcal{A}_\mathcal{E})$, with $\dim(\mathcal{E}) = \sum_{k=1}^m \dim(\mathcal{M}_k) + d$, $d \geq 1$
- (III) a countable set of points $\mathbb{F}^\Sigma \subseteq \mathcal{E}^\Sigma$, satisfying:

$$\mathbb{F}^\Sigma = \begin{cases} \{f^\sigma\}, & \text{if } d = 1 \\ \{f_{(1)}^\sigma, f_{(2)}^\sigma, \dots\} \cong \mathbb{N}, & \text{if } d \geq 1. \end{cases}$$

such that the following topological homeomorphism holds (under inclusion to a subspace)

$$(\mathcal{E}^\Sigma \setminus \mathbb{F}^\Sigma, \mathcal{T}_{\mathcal{E}^\Sigma}|_{\mathcal{E}^\Sigma \setminus \mathbb{F}^\Sigma}) \xrightarrow[\sup]{\cong} (\mathcal{E}, \mathcal{T}_{\mathcal{A}_\mathcal{E}})$$

and $\mathcal{E}$ is an Einstein space of global isometric analytic embedding for $\mathcal{M}_1, \dots, \mathcal{M}_m$.

In addition, $\mathcal{E}^\Sigma$ is **not** even a topological manifold and there does not exists any smoothing resolution of the singularities $\mathbb{F}^\Sigma \subseteq \mathcal{E}^\Sigma$.

Proof. Let $\check{\mathcal{E}} := \mathcal{M}_1 \times \dots \times \mathcal{M}_m \times \mathcal{F}$ be the product of the given semi-Riemannian $\mathcal{M}_k$'s with an analytic manifold $\mathcal{F}$, $\dim(\mathcal{F}) \geq 1$ satisfying the property:

"deleting any countable many points $\mathbb{F}^\Sigma$ from a d -dimensional and a single point $\{f^\sigma\}$ from an 1-dimensional $\mathcal{F}$, the respective complements $\mathcal{F}^{(d)} \setminus \mathbb{F}^\Sigma$ and $\mathcal{F}^{(1)} \setminus \{f^\sigma\}$ are connected."

(If $\dim(\mathcal{F}) = d = 1$, we can choose $\mathcal{F} \cong S^1$ and if $\dim(\mathcal{F}) = d > 1$, $\mathcal{F} \cong \mathbb{R}^d$.)

Its easy to see from the definition of $\check{\mathcal{E}}$ that we can rewrite it as the disjoint union of manifolds

$$\check{\mathcal{E}} = \bigsqcup_{p \in \mathcal{M}_1 \times \dots \times \mathcal{M}_m} \{p\} \times \mathcal{F}$$

For convenience, set

$$n := \sum_{k=1}^m n_k \in \mathbb{N}$$

where $n_k = \dim(\mathcal{M}_k)$, $1 \leq k \leq m$. Then, $\dim(\mathcal{E}) = (\sum_{k=1}^m n_k) + d = n + d$, $d \geq 1$.

Choose a countable set of points

$$\mathbb{F}^\Sigma := \{f_{(1)}^\sigma, f_{(2)}^\sigma, \dots\} \cong \mathbb{N}$$

of $\mathcal{F}$ (if $\dim(\mathcal{F}) > 1$) or a single point $\{f^\sigma\}$ (if $\dim(\mathcal{F}) = 1$). We will prove the general case for $\mathbb{F}^\Sigma$ reducing obviously to the trivial case when $d = 1$.

The previous choice implies a choice of points for each $\{q\} \times \mathcal{F}$ of $\check{\mathcal{E}}$:

$$\{q\} \times \mathbb{F}^\Sigma = \{q\} \times \{f_{(m)}^\sigma\}_{m \in \mathbb{N}} = \{(q, f^\sigma)_{(m)}\}_{m \in \mathbb{N}} \equiv \{f_{q(m)}^\sigma\}_{m \in \mathbb{N}} \cong \mathbb{N}$$

denoted as

$$\mathbb{F}_p^\Sigma := \{f_{p(m)}^\sigma\}_{m \in \mathbb{N}} \subseteq \{p\} \times \mathcal{F}, \quad \forall p \in \mathcal{M}_1 \times \dots \times \mathcal{M}_m$$

Considering the (disjoint) union of all such points in $\check{\mathcal{E}}$

$$\biguplus_{p \in \mathcal{M}_1 \times \dots \times \mathcal{M}_m} \mathbb{F}_p^\Sigma \xrightarrow{\subset} \check{\mathcal{E}} = \biguplus_{p \in \mathcal{M}_1 \times \dots \times \mathcal{M}_m} \{p\} \times \mathcal{F}$$

we may get a more subtle expression of them, as:

$$\mathbb{F}_{(m)}^\Sigma := \{f_{p(m)}^\sigma\}_{p \in \mathcal{M}_1 \times \dots \times \mathcal{M}_m} \subseteq \check{\mathcal{E}} = \mathcal{M}_1 \times \dots \times \mathcal{M}_m \times \mathcal{F}, \quad \forall m \in \mathbb{N}$$

Define now a topological space $\mathcal{E}^\Sigma$ as the quotient space, produced identifying for each m in $\mathbb{N}$ and $\forall p \in \mathcal{M}_1 \times \dots \times \mathcal{M}_m$, all the distinct points $\mathbb{F}_{(m)}^\Sigma$ as one $f_{(m)}^\Sigma$:

$$\mathcal{E}^\Sigma := \check{\mathcal{E}} / \biguplus_{m \in \mathbb{N}} \mathbb{F}_{(m)}^\Sigma = \biguplus_{p \in \mathcal{M}_1 \times \dots \times \mathcal{M}_m} \{\{p\} \times \mathcal{F}\} / \biguplus_{m \in \mathbb{N}} \mathbb{F}_{(m)}^\Sigma$$

with the quotient topology, say $\mathcal{T}_{\mathcal{E}^\Sigma}$, induced by the topologies $\mathcal{T}_{\mathcal{A}_{\mathcal{M}_1 \times \dots \times \mathcal{M}_m}}$ and $\mathcal{T}_{\mathcal{A}_{\mathcal{F}}}$.

In the case of 1-dimensional manifold $\mathcal{F}$, this is exactly the wedge sum of the manifolds $\{p\} \times \mathcal{F}$ with respect to (the 1 term sequence) $\mathbb{F}^\Sigma$ (each f_p^σ is the chosen base point of $\{p\} \times \mathcal{F}$):

$$\mathcal{E}^\Sigma \equiv \bigvee_{\mathbb{F}^\Sigma} \{\{p\} \times \mathcal{F}\} \quad (2.1)$$

Back to the general case, $\check{\mathcal{E}} = \mathcal{M}_1 \times \dots \times \mathcal{M}_m \times \mathcal{F}$ has a natural (product) analytic manifold structure. Deleting all the (countable many) distinct points $\{f_{p(m)}^\sigma\}$, $\forall p \in \mathcal{M}_1 \times \dots \times \mathcal{M}_m$ and $\forall m \in \mathbb{N}$, we obtain an analytic connected manifold structure in the (open in the induced topology) complement:

If we set

$$\mathcal{E} := \check{\mathcal{E}} \setminus \biguplus_{m \in \mathbb{N}} \left(\biguplus_{p \in \mathcal{M}_1 \times \dots \times \mathcal{M}_m} \{f_{p(m)}^\sigma\} \right)$$

then

$$\mathcal{E} = \check{\mathcal{E}} \setminus \bigcup_{p \in \mathcal{M}_1 \times \dots \times \mathcal{M}_m} \mathbb{F}_p^\Sigma = \bigcup_{p \in \mathcal{M}_1 \times \dots \times \mathcal{M}_m} \{p\} \times \{\mathcal{F} \setminus \mathbb{F}_p^\Sigma\}$$

with the obvious C_α^∞ -atlas

$$\mathcal{A}_\mathcal{E} := \mathcal{A}_\mathcal{E}|_\mathcal{E}$$

Deleting the sequence of (identified) points $\{\mathbb{F}_{(m)}^\Sigma\}_{m \in \mathbb{N}}$ of $\mathcal{E}^\Sigma$, we obtain the differential-topological structure of an analytic manifold, $\mathcal{E}^\Sigma \setminus \mathbb{F}^\Sigma$ without any singularities, since:

$$\mathcal{E}^\Sigma \setminus \bigcup_{m \in \mathbb{N}} \mathbb{F}_{(m)}^\Sigma = \{\check{\mathcal{E}} / \bigcup_{m \in \mathbb{N}} \mathbb{F}_{(m)}^\Sigma\} \setminus \bigcup_{m \in \mathbb{N}} \mathbb{F}_{(m)}^\Sigma = \bigcup_{p \in \mathcal{M}_1 \times \dots \times \mathcal{M}_m} \{p\} \times \{\mathcal{F} \setminus \mathbb{F}_p^\Sigma\}$$

that is

$$\mathcal{E}^\Sigma \setminus \bigcup_{m \in \mathbb{N}} \mathbb{F}_{(m)}^\Sigma \equiv \mathcal{E}$$

which proves the analytic structure on $\mathcal{E}^\Sigma \setminus \bigcup_{m \in \mathbb{N}} \mathbb{F}_{(m)}^\Sigma$. Applying now the *embedding theorem* 2.2 for this analytic manifold, we can make it an Einstein global isometric embedding manifold of $\mathcal{M}_1, \dots, \mathcal{M}_m$.

The remaining task is to show that $\mathcal{E}^\Sigma$ is not a manifold. To see this, suppose that taking the wedge sum of 2 distinct fibres $(\{p\} \times \mathcal{F}) \vee_{f_0} (\{q\} \times \mathcal{F})$, there exists a homeomorphism of a domain V_{f_0} of f_0 to $\mathbb{R}^{n+d}$. If this holds, then there must exist a homeomorphism

$$\phi : V \setminus \{f_0\} \xrightarrow{\cong} \phi(V) \setminus \{\phi(f_0)\} \subseteq \mathbb{R}^{n+d}$$

choose an open ball $\mathbb{B}^{n+d}(0, r)$ of maximum radius living into $\phi(V_{f_0})$ and containing $\phi(f_0)$. Then, the following homeomorphism

$$V_{f_0} \cap \phi^{-1}(\mathbb{B}^{n+d}(0, r)) \setminus \{f_0\} \xrightarrow{\cong} \mathbb{B}^{n+d}(0, r) \setminus \{\phi(f_0)\}$$

can not hold, since $\mathbb{B}^{n+d}(0, r) \setminus \{\phi(f_0)\}$ has 1 connected component while $\phi^{-1}(\mathbb{B}^{n+d}(0, r)) \setminus \{f_0\}$ has 2 connected components. □

The mathematical mechanism of topological identification we used to produce the pathological singularities on the Bulk raises the idea that it is not really necessary these "anomalies" to be points: there may exist extended problematic domains of the complement $\mathcal{E}^\Sigma \setminus (\mathcal{M}_1 \times \dots \times \mathcal{M}_m)$. We simply identify distinct homeomorphic domains on the "fibres" $\mathcal{F}$.

This more realistic case may correspond to extended object in the Universe with positive (multi-dimensional) volume.

Theorem 2.2. (*Embedding into Einstein manifold with unsmoothable differential-topological singularities of positive $(d - k)$ -measure*)

Given any finite-many analytic semi-Riemannian manifolds (Branes) $\mathcal{M}_1, \dots, \mathcal{M}_m$ there exist:

(I) *a topological space $(\mathcal{E}^\Sigma, \mathcal{T}_{\mathcal{E}^\Sigma})$*

(II) *an analytic manifold $(\mathcal{E}, \mathcal{A}_{\mathcal{E}})$, with $\dim(\mathcal{E}) = \sum_{k=1}^m \dim(\mathcal{M}_k) + d$, $d \geq 1$*

(III) *countable many closed path connected $\mathcal{E}$ simply connected sets $\mathbb{F}^\Sigma \subseteq \mathcal{E}^\Sigma$, satisfying:*

$$\mathbb{F}^\Sigma = \begin{cases} \{f^\Sigma\} \cong [0, 1], & \text{if } d = 1 \\ \{f_{(1)}^\Sigma, f_{(2)}^\Sigma, \dots\} \cong \mathbb{N} \times \overline{\mathbb{B}^{d-k}(0, 1)}, & \text{if } d \geq 1. \end{cases}$$

($0 \leq k \leq d$) such that the following topological homeomorphism holds (under inclusion to a subspace)

$$(\mathcal{E}^\Sigma \setminus \mathbb{F}^\Sigma, \mathcal{T}_{\mathcal{E}^\Sigma}|_{\mathcal{E}^\Sigma \setminus \mathbb{F}^\Sigma}) \xrightarrow[\supset]{\cong} (\mathcal{E}, \mathcal{T}_{\mathcal{A}_{\mathcal{E}}})$$

*$\mathcal{E}$ is an Einstein space of global isometric analytic embedding for $\mathcal{M}_1, \dots, \mathcal{M}_m$. In addition, $\mathcal{E}^\Sigma$ is **not** a topological manifold.*

Proof. The proof is a replica of the previous result. The only fact that needs to be noted is that extracting from a closed (abstract) curve a closed (image of) an interval, (e.g. a closed arc from S^1) or a closed (image of a) $(d - k)$ -ball from $\mathcal{F}$ (choosing e.g. $\mathbb{R}^d$) for $\dim(\mathcal{F}) = d$, we obtain open submanifolds in the analytic subspace structure.

In addition, the inability of resolution of the singularities, owes to the fact that choosing from each of the $\mathcal{F}$'s a (diffeomorphic image of a) closed ball,

the quotient space when they are all identified is not a manifold, by the same topological argument as in 2.1, extracting a closed inversed image of ball instead of a point.

Countability of these "extended anomalies" can be easily shown, since choosing for the (diffeomorphic images of) balls, $\text{diam}(\overline{\mathbb{B}^{d-k}(0, r)}) = 2r < 2$, we can embed in $\mathbb{R}^d$ countable many balls, centered at the points of $\mathbb{N}^d$. If $k \neq 0$, it suffices to reduce embedding of balls to a $d - k$ -dimensional linear subspace of $\mathbb{R}^d$. $\square$

Acknowledgement. *The Author is profoundly indebted to professor Lorenz Magaard at University of Honolulu, Hawaii for his interest in the present work, P. Krikelis & A. Kartsaklis at University of Athens for helpful comments & technical support and Ed. Anderson for discussions on the physical aspects of the embedding results.*

References

- [1] Glen E. Bredon, *Topology and Geometry*, Springer, 1997
- [2] J. E. Campbell, *A Course of Differential Geometry*, Oxford, Clarendon Press, 1923
- [3] Do Carmo, M.P., *Riemannian Geometry*, Birkhäuser, 1992
- [4] L. Pf. Eisenhart, *Riemannian Geometry*, Princeton University Press, 1964
- [5] O. Gron and S. Hervik, *Einstein's General Theory of Relativity* (e-book), <http://www.fys.uio.no/~sigbjorh/gravity.pdf>, 2004
- [6] A. Hatcher, *Algebraic Topology*, <http://www.math.cornell.edu/~hatcher>, (e-book), 2000
- [7] S. W. Hawking & G. F. R. Ellis *The Large scale structure of spacetime*, Cambridge University Press, 1973
- [8] L. Magaard, *Zur Einbettung Riemannscher Raume in Einstein - Raume und Konfor - Euclidische Raume*, PhD thesis, (Kiel), 1963

- [9] J. R. Munkres, *Topology, a first course*, Prentice-Hall, Inc., Englewood Cliffs, New Jersey
- [10] R. Narasimhan, *Analysis on Real and Complex Manifolds*, Masson and Cie, Paris, North Holland, 1968
- [11] R. Penrose, *Techniques of Differential Topology in Relativity*, SIAM Publ. Philadelphia, 1972
- [12] R. K. Sachs and H. Wu, *General Relativity for Mathematicians*, Springer-Verlang, 1977
- [13] N. Steenrod, *The topology of fibre Bundles*, Princeton University Press, 1972
- [14] H. Stefani, *General Relativity*, 2nd Edition, Cambridge University Press, 1990
- [15] P. S. Wesson, *Space-Time-Matter, Modern Kaluza-Klein Theory*, World Scientific, 2000
- (Articles)
- [16] E. Anderson, J. E. Lidsey, *Embeddings in NonVacuum Spacetimes*, Class.Quant.Grav. 18 (2001) 4831-4844, ArXiv: gr-qc/0106090
- [17] F. Dahia, C. Romero, *The embedding of spacetime in 5 dimensions: An extension of Campbell-Magaard theorem*, J.Math.Phys. 43 (2002) 5804-5814, ArXiv: gr-qc/0109076
- [18] F. Dahia, C. Romero, *On the embedding of branes in five-dimensional spaces*, Class.Quant.Grav. (2004) 21, No 4, 927-933, ArXiv: gr-qc/0308056
- [19] A. Davidson, $\Lambda = 0$ cosmology of a brane-like universe, Class. Quantum Grav. 16 (1999) 653-659
- [20] C. Deffayet, *Cosmology of Brane-Induced Gravity Models*, International Journal of Theoretical Physics, Vol. 41, No. 11, November 2002
<http://www.kluweronline.com/article>

- [21] M. Gogberashvili, *Brane-Universe in Six Dimensions with Two Times*, Phys.Lett. B484 (2000) 124-128, ArXiv: hep-ph/0001109
- [22] M. Gogberashvili, *Embedding of the Brane into Six Dimensions*, J.Math.Phys. 43 (2002) 4886 - 4888, ArXiv: gr-qc/0202061
- [23] M. Gogberashvili, *Brane Gravity from Bulk Vector Field*, Phys.Lett. B553 (2003) 284-288, ArXiv: hep-th/0203090
- [24] H. Grauert, *On Levi's problem and the imbedding of real analytic manifolds*, Annals of Mathematics, 68 (47-82)
- [25] E. Gravanis, S. Willison, *Israel conditions for the Gauss-Bonnet theory and the Friedmann equation on the brane universe*, ArXiv: hep-th/0209076
- [26] N. I. Katzourakis "The Global Embedding Problem of Semi- Riemannian into Einstein Manifolds", ArXiv: math-ph/0407067
- [27] N. I. Katzourakis "On the Global Embedding of Spacetime into Singular $\mathcal{E}^\Sigma$ Einstein Manifolds: Wormholes", ArXiv: math.DG/0503129
- [28] N. I. Katzourakis "Global Embedding of Analytic Branes into Einstein MD Bulk Cosmology", ArXiv: math.DG/0411630
- [29] D. Langlois, *Brane cosmology: an introduction*, Prog.Theor.Phys.Suppl. 148 (2003) 181-212, ArXiv: hep-th/0209261
- [30] D. Langlois, *Gravitation and Cosmology in Brane-Worlds*, (Invited plenary talk at the GR17 conference, Dublin, Ireland, July 2004), ArXiv: gr-qc/0410129
- [31] J. E. Lidsey, C. Romero, R. Tavakol, S. Rippl, *On Applications of Campbell's Embedding Theorem*, Class.Quant.Grav. 14 (1997) 865-879, ArXiv: gr-qc/9907040
- [32] J. E. Lidsey, *The Embedding of Superstring Backgrounds in Einstein Gravity*, Phys.Lett. B417 (1998) 33, ArXiv: hep-th/9907095

- [33] R. Maartens, *Geometry and Dynamics of the Brane-World*, Reference Frames and Gravitomagnetism, ed. J Pascual-Sanchez et al. (World Sci., 2001), p. 93-119, ArXiv: gr-qc/0101059
- [34] R. Maartens, *Brane-World Gravity*, Living Rev. Relativity, 7, (2004), 7, <http://www.livingreviews.org/lrr-2004-7>
- [35] M. Pavisic, *Resource Letter on Geometrical Results for Embeddings and Branes*, arXiv: gr-qc/0010045
- [36] A. Perez-Lorenzana, *An Introduction to the Brane World*, ArXiv: hep-ph/0406279
- [37] F. Quevedo, *Lectures on String/Brane Cosmology*, arXiv: hep-th/0210292
- [38] L. Randall, R. Sundrum, *A Large Mass Hierarchy from a Small Extra Dimension*, Phys.Rev.Lett. 83 (1999) 3370-3373, ArXiv: hep-ph/9905221
- [39] L. Randall, R. Sundrum, *An Alternative to Compactification*, Phys.Rev.Lett. 83 (1999) 4690-4693, ArXiv: hep-th/9906064
- [40] S. S. Seahra, *Physics in Higher-Dimensional Manifolds*, <http://etd.uwaterloo.ca/etd/ssseahra2003.pdf>
- [41] S. S. Seahra, P. S. Wesson, *Application of the Campbell- Magaard theorem to higher-dimensional physics*, Class.Quant.Grav. 20 (2003) 1321-1340, ArXiv: gr-qc/0302015
- [42] T. Shiromizu, K. Koyama, *Low energy effective theory for two branes system - Covariant curvature formulation*, Phys.Rev. D67 (2003) 084022, ArXiv: hep-th/0210066
- [43] D. A. Steer, M. F. Parry, *Brane Cosmology, Varying Speed of Light and Inflation in Models With One or More Extra Dimensions*, Int.J.Th.Phys., Vol. 41, No. 11, Nov. 2002, <http://www.kluweronline.com/article>
- [44] D. Wands, *Brane-world cosmology*, <http://www.tech.port.ac.uk/staffweb/wandsd/powerpoint/sussex.pdf>

- [45] H. Whitney, *Differentiable Manifolds*, Annals of Mathematics, 37 (645-680), 1936